

**NORM INEQUALITY FOR THE CLASS OF SELF-ADJOINT  
ABSOLUTE VALUE GENERALIZED DERIVATIONS**

**Danko R. Jocić**

**Abstract.** We prove that for all  $0 \leq \alpha \leq 2/3$

$$\||A|^\alpha X - X|B|^\alpha\| \leq 2^{2-\alpha} \|X\|^{1-\alpha} \|AX - XB\|^\alpha,$$

for all bounded Hilbert space operators  $A = A^*$ ,  $B = B^*$  and  $X$ , as well as

$$\||A|^\alpha - |B|^\alpha\| \leq 2^{2-\alpha} \|A - B\|^\alpha,$$

for arbitrary bounded  $A$  and  $B$ .

Let  $H$  be a complex, infinite dimensional Hilbert space,  $B(H)$  the algebra of all bounded linear operators on  $H$  and let  $\|\cdot\|$  stands for the norm in  $B(H)$ . The following theorem compares a class of the absolute value generalized derivations on  $B(H)$ , induced by a pair of self-adjoint operators.

**THEOREM 1.** For all  $0 \leq \alpha \leq 2/3$  we have

$$\||A|^\alpha X - X|B|^\alpha\| \leq 2^{2-\alpha} \|X\|^{1-\alpha} \|AX - XB\|^\alpha,$$

for bounded Hilbert space operators  $A = A^*$ ,  $B = B^*$  and  $X$ .

*Proof.* Let  $A = U|A|$  and  $B = V|B|$  be polar decompositions of  $A$  and  $B$ , with unitary  $U = U^*$  and  $V = V^*$ ,  $|A| = \sqrt{A^*A}$  and  $|B| = \sqrt{B^*B}$ . Thus

$$\begin{aligned} \||A|^\alpha X - X|B|^\alpha\| &= \\ &\|U|A|^{\frac{\alpha}{2}} (U|A|^{\frac{\alpha}{2}} X - XV|B|^{\frac{\alpha}{2}}) + (U|A|^{\frac{\alpha}{2}} X - XV|B|^{\frac{\alpha}{2}}) V|B|^{\frac{\alpha}{2}}\| \\ &\leq 2\|U|A|^{\frac{\alpha}{2}} X - XV|B|^{\frac{\alpha}{2}}\|^{\frac{1-\alpha}{1-\alpha/2}} \times \\ &\left\| \frac{|A|^{1-\frac{\alpha}{2}} (U|A|^{\frac{\alpha}{2}} X - XV|B|^{\frac{\alpha}{2}}) + (U|A|^{\frac{\alpha}{2}} X - XV|B|^{\frac{\alpha}{2}}) |B|^{1-\frac{\alpha}{2}}}{2} \right\|^{\frac{\alpha/2}{1-\alpha/2}}, \end{aligned} \tag{1}$$

*AMS Subject Classification:* 47 A 30, 47 B 05, 47 B 10, 47 B 15

*Keywords and phrases:* singular values, three line theorem for operators, unitarily invariant norms

Communicated at the 4th Symposium on Mathematical Analysis and Its Applications, Arandelovac 1997.

by Corollary 2.2 of [2] applied to  $U|A|^{\frac{\alpha}{2}}X - XV|B|^{\frac{\alpha}{2}}$  instead of  $X$  and  $r = \frac{2-\alpha}{\alpha} \geq 2$ . As  $\alpha/2 \leq 1/3$ , then an application of Theorem 3.1 of [1] for  $p = 2/\alpha \geq 3$  shows that

$$\|U|A|^{\frac{\alpha}{2}}X - XV|B|^{\frac{\alpha}{2}}\| \leq \|2X\|^{1-\frac{\alpha}{2}} \|AX - XB\|^{\frac{\alpha}{2}}. \quad (2)$$

Also, we have

$$\begin{aligned} & \| |A|^{1-\frac{\alpha}{2}}(U|A|^{\frac{\alpha}{2}}X - XV|B|^{\frac{\alpha}{2}}) + (U|A|^{\frac{\alpha}{2}}X - XV|B|^{\frac{\alpha}{2}})|B|^{1-\frac{\alpha}{2}} \|/2 \\ &= \|AX - XB + (U|A|^{\frac{\alpha}{2}}X|B|^{1-\frac{\alpha}{2}} - |A|^{1-\frac{\alpha}{2}}XV|B|^{\frac{\alpha}{2}})\|/2 \\ &\leq \|AX - XB\|, \end{aligned} \quad (3)$$

by Lemma 3.2 of [1] applied for  $p = 1$  and  $s = \frac{\alpha}{2}$ . Now, according to (2) and (3), (1) finally gives

$$\begin{aligned} \| |A|^{\alpha}X - X|B|^{\alpha} \| &\leq 2\|2X\|^{(1-\frac{\alpha}{2})\frac{1-\alpha}{1-\alpha/2}} \|AX - XB\|^{\frac{\alpha}{2}\frac{1-\alpha}{1-\alpha/2} + \frac{\alpha/2}{1-\alpha/2}} \\ &= 2^{2-\alpha} \|X\|^{1-\alpha} \|AX - XB\|^{\alpha}. \quad \blacksquare \end{aligned} \quad (4)$$

This theorem also enables us to derive the following perturbation result for a class of the absolute value map in  $B(H)$ .

**THEOREM 2.** *For all  $0 \leq \alpha \leq 2/3$  we have*

$$\| |A|^{\alpha} - |B|^{\alpha} \| \leq 2^{2-\alpha} \|A - B\|^{\alpha}, \quad (5)$$

for arbitrary bounded Hilbert space operators  $A$  and  $B$ .

*Proof.* Define  $C = \begin{bmatrix} 0 & A^* \\ A & 0 \end{bmatrix}$  and  $D = \begin{bmatrix} 0 & B^* \\ B & 0 \end{bmatrix}$  as operators acting on  $H \oplus H$ . A straightforward calculation shows that  $C = C^*$ ,  $D = D^*$ ,  $|C|^{\alpha} = \begin{bmatrix} |A|^{\alpha} & 0 \\ 0 & |A^*|^{\alpha} \end{bmatrix}$  and  $|D|^{\alpha} = \begin{bmatrix} |B|^{\alpha} & 0 \\ 0 & |B^*|^{\alpha} \end{bmatrix}$ . Also

$$\|C - D\| = \max\{\|A - B\|, \|A^* - B^*\|\} = \|A - B\|$$

and

$$\| |A|^{\alpha} - |B|^{\alpha} \| \leq \max\{\| |A|^{\alpha} - |B|^{\alpha} \|, \| |A^*|^{\alpha} - |B^*|^{\alpha} \| \} = \| |C|^{\alpha} - |D|^{\alpha} \|.$$

An application of the preceding theorem to self-adjoint  $C$  and  $D$  and  $X = I$  gives

$$\| |A|^{\alpha} - |B|^{\alpha} \| \leq \| |C|^{\alpha} - |D|^{\alpha} \| \leq 2^{2-\alpha} \|C - D\|^{\alpha} = 2^{2-\alpha} \|A - B\|^{\alpha},$$

completing the proof.  $\blacksquare$

#### REFERENCES

- [1] Danko R. Jocić, *Norm inequalities for self-adjoint derivations*, J. Functional Analysis **145** (1997), 24-34.
- [2] Danko R. Jocić, *Cauchy-Schwartz and means inequalities for elementary operators into norm ideals*, (to appear in the Proc. Amer. Math. Soc.)

(received 22.09.1997.)

University of Belgrade, Faculty of Mathematics, Studentski trg 16, P.P. 550, 11000 Belgrade, Yugoslavia

E-mail: Danko@rlab.matf.bg.ac.yu and jocić@matf.bg.ac.yu